

FORCING MECHANISMS FOR DESTRUCTIVE SEICHES IN THE BALEARIC ISLANDS

David C. Chapman
Woods Hole Oceanographic Institution
Woods Hole, MA 02543
phone: (508) 289-2792 fax: (508) 457-2181 email: dchapman@whoi.edu

Graham S. Giese
Woods Hole Oceanographic Institution
Woods Hole, MA 02543
phone: (508) 289-2297 fax: (508) 457-2187 email: ggiese@whoi.edu

Award #: N0001492J1200

LONG-TERM GOALS

Our long-term goals are to increase understanding of the role played by internal waves in coastal seiche generation globally and to apply this knowledge to improve hazardous seiche prediction.

OBJECTIVES

The objectives of the present effort are to investigate the forcing mechanisms for the large-amplitude, destructive seiching events that occur at Ciutadella Harbor in the Balearic Islands (western Mediterranean) and, based on the results, to develop methodologies for providing advance warning of those events.

APPROACH

We developed a cooperative field study in collaboration with scientists at the University of the Balearic Islands (UBI) to gather data from Ciutadella Harbor and vicinity to test hypotheses concerning the forcing of the destructive seiching events (e.g., forcing is primarily by meteorological processes; forcing is primarily by internal waves). A sea level and water temperature recorder was established in the harbor, a mooring with water temperature and pressure sensors was established offshore in Menorca Passage, and wind data were obtained from the Spanish meteorological service.

WORK COMPLETED

We have collected one-minute sea level and water temperature data at Ciutadella Harbor for 1123 days during 1992-1996, water temperature and pressure data from the offshore mooring for four months during 1994, and surface wind data for the same four-month period. Analysis of these data was completed in 1997.

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Forcing Mechanisms for Destructive Seiches in the Balearic Islands				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Woods Hole Oceanographic Institution, Woods Hole, MA, 02543				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a REPORT unclassified	b ABSTRACT unclassified	c THIS PAGE unclassified			

RESULTS

Ten-minute (resonant) oscillations of Ciutadella Harbor were almost always discernible from the sea-level data, but seiches larger than about 0.5 m had a pronounced seasonal distribution closely correlated with the seasonal distribution of surface water (upper 50 m) stratification (fig. 1). Extreme seiching events (daily mean activity greater than 50 percent of the total range for all 1123 days) occurred only four times during the observation period, once each during 1992, 1993, 1994 and 1996. Maximum seiche oscillations during these events ranged between 1 and 2 meters.

The four extreme events each occurred under unusual and remarkably similar tidal conditions, near-apogean neap tides. The four-month period for which we had wind data included the 1994 event. The steadiest episode of strong southeasterly wind of the entire four-month period immediately preceded the extreme seiching event. Ciutadella fishermen have often reported a similar relationship between southeasterly wind and destructive seiche events.

IMPACT/APPLICATIONS

Previously, it was thought that the role of internal waves in large-amplitude seiche generation was restricted to those cases in which seiches are forced by tide-generated internal waves. The new results provide evidence that both tides and internal waves play an important role even when the initial forcing is provided by atmospheric processes. In such cases, application of this improved understanding should produce improved predictions of destructive events.

TRANSITIONS

Prior to the 1998 "seiche season" we expect to be able to provide our colleagues at UBI dates of the approximately 6 periods in 1998 when water stratification and tidal conditions would "allow" for extreme seiching events. We would expect such an event on any of those dates if strong and sustained southeasterly winds also occur.

RELATED PROJECTS

Julio Candela and Richard Limeburner (Woods Hole Oceanographic Institution) are studying flows through major passages of the western Mediterranean. Clint Winant and colleagues (Scripps Institution of Oceanography) have obtained interesting measurements of shoreward propagating internal waves over the shelf off southern California.




Figure 1: Relative daily sea-level variance (left axis) at Ciutadella Harbor for periods between 9 and 12 minutes ('x') plotted against year day. This plot represents data for 1123 days. Dashed horizontal line on left side indicates value of 50% of the total range of daily variance and separates the four days of exceptional seiche activity from the remainder. At relative variance values of approximately 3, seiches with ranges as large as 0.5 m may occur; at values of about 10, seiche ranges as large as 1 m may occur. Circles connected by solid lines indicate the mean monthly harbor surface water temperature during the study period and refer to the vertical axis on the right side of the figure. Since offshore water at a depth of 50 m maintains a relatively stable temperature of approximately 13 degrees C, these data indicate monthly changes in thermal stratification.

